

INTEGRAL and Swift observations of the hard X-ray transient MAXI J1828-249

E. Filippova^{1,2}, E. Bozzo¹, and C. Ferrigno¹

¹ ISDC, University of Geneva, Ch. d'Ecogia 16, 1290 Versoix, Switzerland

² Space Research Institute, Moscow, 117997, Russia; e-mail: Ekaterina.Filippova@unige.ch

Received / Accepted

ABSTRACT

In this paper we report on the observations performed with *INTEGRAL* and *Swift* of the first outburst detected from the hard X-ray transient MAXI J1828-249. During the first about two days of the outburst, the source was observed by MAXI to undergo a very rapid transition from a hard to a softer spectral state. While the hard state was not efficiently monitored because the transition occurred so rapidly, the evolution of the source outburst in the softer state was covered quasi-simultaneously in a broad energy range (0.6-150 keV) by the instruments on-board *INTEGRAL* and *Swift*. During these observations, the spectra measured from the source displayed both a prominent thermal emission with temperature $kT \sim 0.7$ keV and a power-law hard component with a photon index $\Gamma \sim 2.2$ extending to 200 keV. The properties of the source in the X-ray domain are reminiscent of those displayed by black hole transients during the soft intermediate state, which supports the association of MAXI J1828-249 with this class of objects.

Key words. methods:observational - X-rays:binaries - X-rays:individuals: MAXI J1828-249

1. Introduction

MAXI J1828-249 was discovered in outburst on 15 October 2013 (56580.91 MJD) by MAXI/GSC (Nakahira et al. 2013). Shortly after the discovery, the source was detected at higher energies by the hard X-ray imager ISGRI on-board INTEGRAL (Filippova et al. 2013). At discovery the flux recorded from the source was 93 ± 9 mCrab in the 4-10 keV energy band and 45 ± 2 mCrab (48 ± 2 mCrab) in the 20-40 keV (40-80 keV) energy band. Follow-up observations carried out on 56581.6 MJD with the narrow field instrument on-board *Swift*/XRT provided the best measured position of the source at RA=277.2427, Dec=-25.0304 (J2000) with an estimated uncertainty of 3.6 arcsec at 90 c.l. (Kennea et al. 2013a). The improved position permitted us to identify the UV and IR counterparts of the source. The estimated magnitude of the UV source was 18.64 ± 0.04 (stat) ± 0.03 (sys) (in AB system, without correction for interstellar reddening; Kennea et al. 2013b). The AB magnitudes of the optical/IR source were $g' = 17.2 \pm 0.1$, $r' = 16.9 \pm 0.1$, $i' = 16.9 \pm 0.1$, $z' = 16.8 \pm 0.1$, $J = 16.8 \pm 0.1$, $H = 16.9 \pm 0.1$, $K = 17.2 \pm 0.2$ (not corrected for the interstellar reddening). The derived spectrum of optical/IR source might be interpreted as arising because of an accretion disk (Rau et al. 2013). Radio observations carried out about two days after the onset of the outburst did not reveal significant emission from the source (the 3σ upper limit on its flux was $75 \mu\text{Jy}$ at 5.5 GHz and $57 \mu\text{Jy}$ at 9.0 GHz; Miller-Jones et al. 2013).

In this paper we report on all *INTEGRAL* and *Swift* data available to our group collected during the outburst of MAXI J1828-249 from 56580 to 56607 MJD.

2. INTEGRAL data

MAXI J1828-249 was observed by IBIS/ISGRI (Lebrun et al. 2003; Ubertini et al. 2003) from 56580 MJD to 56594 MJD,




Fig. 1. ISGRI mosaic of the FOV around MAXI J1828-249 (20-100 keV). The mosaic has been obtained from data collected during revolution 1344.

that is during satellite revolutions 1344 to 1348. Revolution 1348 closed the seasonal visibility window of *INTEGRAL* in the direction of the source and no other data were collected after 56594 MJD (see Table 1). The source was generally observed at relatively high off-axis angles and was included in the FOV of the two JEM-X units (Lund et al. 2003) for only a relatively small fraction of the total observational time (revolutions 1346 to 1348). The analysis of *INTEGRAL* data was carried out by the OSA10.0 distributed by the ISDC (Courvoisier et al. 2003). We show in Fig. 1 the ISGRI mosaic of the field of view (FOV) around the source. In this mosaic (total effective exposure time 137 ks) the source is detected with a significance of 41σ in the 20-100 keV energy band. In the JEM-X mosaic (effective exposure time 6 ks), the source was detected at a significance of 13σ in the 3-35 keV energy band. To perform a broad-band spectral analysis of the X-ray emission recorded from the source, we extracted the ISGRI and JEM-X spectra during time intervals quasi-simultaneous with *Swift* data (see Table 1).

3. Swift data

Swift observations were carried out immediately after the discovery of the source and covered about 30 days of the outburst (Kennea et al. 2013a,b). The first two pointings (ID 00032993001, 00032994001) were collected in photon-counting mode (PC; time resolution 2.5 s) and were severely affected by pile-up due to the high flux of the source (Kennea et al. 2013a). We therefore discarded these data for the following analysis. All the remaining data were collected in window-timing mode (WT; time resolution 1.8 ms, see Table 2). XRT data were processed with the `XRTPIPELINE` (v.0.12.6) and were analyzed using standard procedures (Burrows et al. 2005). We selected only event grades 0-2 and adopted the latest response files available (v.014). Exposure maps were created through the `XRTPEXMAP` task and were used to generate ARF files. The source light curves in each observation were corrected for all instrumental issues with the task `XRTLCCORR`. All source event lists were barycentered by using the `BARYCORR` tool. In all XRT observations the source was detected at count-rates 150-250 cts/s in energy band 0.3-10 keV, depending also on the position of the source on the detector and the presence of bad columns¹. We followed the technique described in Romano et al. (2006) to search for pile-up in the XRT observations, and found that excluding the two innermost pixels of the extraction region centered on the source was sufficient to correct for this problem in all pointings. Correction for pile-up was not needed for pointing 8, as in this case a bad column passing through the source already decreased the count-rate below the level at which pile-up is a problem. All XRT spectra were grouped to have at least 100 photons per energy bin and were fit with the χ^2 -statistic in *XSPEC*. Uncertainties throughout the paper are given at 90% c.l. if not stated otherwise.

4. Results

The XRT and ISGRI light curves of the source covering the entire observational campaign reported in this paper are shown in Fig. 4. The XRT light curve shows a noticeable increase of the source X-ray flux (0.3-10 keV) during the first five days of the outburst with a gradual decrease afterwards. In the ISGRI energy band (20-100 keV) no significant flux evolution is detected (though with large uncertainties). This behavior agrees with that found in the publicly available MAXI data of the source².

We first extracted all available XRT spectra of the source and fit them with a simple absorbed power-law model. In all cases these fits gave unacceptable results ($\chi^2_{\text{red}} \gg 1$) and showed a clear excesses in the residuals at energies $\lesssim 2$ -3 keV. Data points at energies < 0.6 keV were discarded from the analysis to avoid strong instrumental residuals around the O-K edge. We found that a model comprising an absorbed multi-temperature disk black body (`DISKBB` in *XSPEC*) and a power law could be used to describe all XRT spectra in the energy range 0.6-10 keV equally well. In all cases we obtained an absorption column density consistent with the Galactic value expected in the direction of the source ($\sim 0.3 \times 10^{22} \text{ cm}^{-2}$, Dickey & Lockman 1990), an inner disk radius (temperature) of $R_{\text{bb}} \sim 25$ km ($kT_{\text{bb}} \sim 0.7$ keV) and a power-law photon index of $\Gamma \sim 3$. When the *INTEGRAL* data were also used to perform a broad-band fit, the higher energy emission of the source was characterized by a significantly harder power-law spectrum ($\Gamma=2.2$) than that derived from the XRT spectra. We found that this discrepancy arises from instrumental residuals in the XRT spectra at energies 1.5-2.5 keV,




Fig. 2. MAXI J1828-249 lightcurve during the outburst as observed by *Swift*/XRT (0.3-10 keV, upper panel) and *INTEGRAL*/ISGRI (20-100 keV, lower panel). The time bin is 1 ks for XRT and 16 ks for ISGRI.

corresponding to the Au – M_V and Si – K edges³. These residuals are tackled by the χ^2 minimization routine in *XSPEC* mainly by adjusting the power-law photon index (poorly determined in the fits to the XRT spectra alone because of the limited energy coverage of the instrument). The instrumental residuals in XRT become very evident when all data are summed to extract an average spectrum (see Fig. 3). Following the suggestion in the latest available XRT calibration document, we tried to account for these residuals in the fits to the XRT data alone by fixing $\Gamma = 2.2$ (as determined from the average broad-band spectrum) and using the `GAIN` correction in *XSPEC*. The results of the broad-band spectral analysis obtained with this method are summarized in Table 1 (normalization constants were introduced in all broad-band fits to account for the variability of the source and intercalibration between the different instruments). We indicated in all cases the measured correction to the energy slope (SI) and shift (G) of the XRT spectra. For consistency, the gain correction was also applied during the fits to the XRT spectra alone, as reported in Table 2 (in several cases the values of SI and G were poorly constrained and did not significantly affect the fits because of the relatively low statistics of the data). The hard X-ray emission from the source is clearly detected up to ~ 200 keV.

The timing analysis of the *Swift*/XRT data was carried out with a technique similar to that reported in Ferrigno et al. (2012). We first searched for timing features in the event files extracted from each of the XRT pointing, but no statistically significant features were found. We therefore computed the average power spectral density (PSD) of the source by performing a Fourier transform on its light curves (time bin of 1 s) and averaging on intervals of 512 s (corresponding to roughly 1-2 time intervals per observation). The PSD were well fit ($\chi^2_{\text{red}}/\text{d.o.f.} = 1.3/33$) by a zero-centered Lorentzian plus white noise. The latter was added as a constant C to the fit and we measured in Leahy normalization $C = 2.17 \pm 0.07$. The source PSD is shown in Fig. 4. We estimated the rms variability of the source from the zero-centered Lorentzian after subtracting the contribution of the white noise and obtained $\text{rms} = 0.90^{+0.11}_{-0.13}\%$ and $\nu_0 = 14^{+8}_{-4}$ mHz. Because this average analysis showed that the aperiodic noise is more prominent in the 4-50 mHz frequency range, we used this interval to compute the fractional rms of each observation sepa-

¹ <http://www.swift.ac.uk/analysis/xrt/exposuremaps.php>

² <http://maxi.riken.jp/top/index.php?cid=1&jname=J1828-250>

³ http://www.swift.ac.uk/analysis/xrt/files/SWIFT-XRT-CALDB-09_v1.7.pdf




Fig. 3. Averaged MAXIJ1828-249 broad-band spectrum obtained by integrating over all available *Swift*/XRT (black), JEM-X (red) and ISGRI (green) data from MJD 56580 to MJD 56594 ("total" spectrum in Table 1). The best-fit model is shown with a solid line. The two dashed lines represent the spectral components (the multi-temperature disk black-body and the power-law). Residuals from the best fit are reported in the middle (no gain correction applied) and bottom panel (after gain correction). Some instrumental residuals are still visible at energies 1.5-2.5 keV also after the gain correction but their significance is strongly reduced.

Table 1. Results of the broad-band spectral analysis.

Data ¹	Exposures (ks)			N_H ²	Γ	kT_{bb} keV	R_{disk} ³ (km)	C_{JEM-X}	C_{ISGRI}	Flux ⁴		SI ⁵	G ⁵ eV	$\chi^2/d.o.f.$
	XRT	JEM-X	ISGRI							0.6-10 keV	20-100 keV			
1344+01	1.0	-	26	0.19±0.02	2.0±0.16	0.68±0.02	28.4±1.8	-	0.6±0.2	4.9	0.6	0.96±0.01	37±25	1.2/283
1344+03	1.0	-	8	0.32±0.08	2.68±0.24	0.73±0.03	22.7±2.5	-	1.9±0.9	6.1	0.7	0.99±0.02	0±30	1.2/331
1345+06	1.0	-	30	0.2±0.02	2.1±0.2	0.76±0.01	25.3±1.7	-	0.7±0.2	6.1	0.6	0.97±0.01	35±18	1.2/338
1346+07	0.9	-	9	0.18±0.03	1.86±0.24	0.74±0.02	27.8±1.7	-	0.5 ^{+0.2} _{-0.1}	6.9	0.8	0.92±0.02	19 ⁺³² ₋₂₇	1.0/310
1346+08	0.2	1.7	17	0.21±0.04	2.1±0.2	0.7±0.04	22.7±3.3	2.3 ^{+0.9} _{-0.7}	1.4 ^{+0.8} _{-0.4}	3.5	0.7	1.03±0.03	25 ⁺³⁰ ₋₅₀	1.0/121
1347+11	0.9	3.4	30	0.25±0.05	2.3±0.3	0.68±0.02	29±3	0.5±0.2	0.5 ^{+0.4} _{-0.2}	5.7	0.4	0.98±0.02	15±20	1.2/251
1348+12	1.05	3.4	40	0.2±0.02	2.2±0.2	0.72±0.01	25.6±1.4	0.7±0.1	0.63±0.03	6.1	0.6	0.95±0.02	45 ⁺²⁰ ₋₂₂	1.1/211
Total	10.5	8.6	160	0.22±0.01	2.2±0.1	0.72±0.04	25.4±0.5	0.7±0.1	0.8±0.1	5.9	0.6	0.97±0.03	24±6	1.4/614

Notes. ¹Indicates the INTEGRAL revolution + the latest two digits XX of the Swift observation ID 000329970XX. ²In units of $\times 10^{22} \text{cm}^{-2}$. ³The radius of the disk is calculated assuming $\cos \theta = 0$ and a distance to the system of 8 kpc. ⁴In units of $\times 10^{-9} \text{erg s}^{-1} \text{cm}^{-2}$. ⁵G (SI) is the energy shift (slope) of the gain correction derived from the spectral fit in Xspec.

rately. The results are reported in Fig. 5. The uncertainties on the rms of each observation was obtained from bootstrapping simulations with 1000 realizations (for a detailed description of the method see Ferrigno et al. 2012). The source exhibited an rms $\leq 10\%$ in all XRT pointings and no particular trend is found when the rms is plotted as a function of the total source count-rate (see Fig. 5).

Owing to the relatively low statistics of the *INTEGRAL* data, we did not attempt a timing analysis of the hard X-ray emission from MAXIJ1828-249.

5. Discussion

MAXIJ1828-249 was discovered in outburst on 2013 October 15 by the MAXI/GSC instrument and apparently underwent a

Table 2. *Swift* data spectral fits.

ObsID ¹	Start Time MJD	Stop Time MJD	Exposure ks	N_H ($\times 10^{22} \text{ cm}^{-2}$)	Photon index ³ Γ	kT_{bb} keV	R_{disk}^2 km	Flux (0.6 – 10 keV) $10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2}$	SI ⁵	G ⁵ eV	$\chi^2/\text{d.o.f}$
1	56581.831	56581.860	0.97	0.21 \pm 0.02	2.2	0.67 \pm 0.02	28.1 $^{+1.6}_{-1.7}$	4.9	0.96 \pm 0.01	29 $^{+26}_{-20}$	1.1/275
2	56582.720	56582.791	0.98	0.21 \pm 0.03	2.2	0.72 \pm 0.02	26.4 $^{+1.7}_{-1.2}$	6.4	0.96 \pm 0.01	45 $^{+16}_{-28}$	1.0/311
3	56583.244	56583.256	0.95	0.22 \pm 0.02	2.2	0.71 \pm 0.01	26.5 \pm 1.2	6.2	0.97 \pm 0.01	20 $^{+21}_{-21}$	1.2/324
4 ⁴	56583.509	56583.517	0.68	0.23 \pm 0.03	2.2	0.70 \pm 0.03	28.2 $^{+3.4}_{-2.1}$	6.5	0.97 \pm 0.02	12 $^{+45}_{-33}$	1.1/250
4 ⁴	56584.579	56584.586	0.59	0.23 \pm 0.03	2.2	0.73 \pm 0.02	29.6 $^{+2.1}_{-2.3}$	6.8	0.97 \pm 0.02	7 $^{+35}_{-27}$	1.1/265
5	56585.039	56585.053	1.20	0.21 \pm 0.02	2.2	0.75 \pm 0.01	27.8 $^{+1.9}_{-1.3}$	6.8	1.01 \pm 0.01	0 $^{+18}_{-28}$	1.2/352
6	56586.185	56586.196	1.01	0.20 \pm 0.01	2.2	0.78 \pm 0.01	24.9 $^{+1.1}_{-1.0}$	6.1	0.97 \pm 0.01	35 \pm 16	1.1/330
7	56587.378	56587.389	0.94	0.22 \pm 0.02	2.2	0.74 \pm 0.02	27.3 \pm 2.0	6.7	0.98 \pm 0.02	0 $^{+32}_{-17}$	1.0/303
8	56588.973	56588.975	0.18	0.22 \pm 0.03	2.2	0.70 \pm 0.04	23.0 \pm 3.0	3.5	1.03 \pm 0.04	0 $^{+35}_{-43}$	0.97/107
11 ⁴	56591.308	56591.318	0.90	0.23 \pm 0.02	2.2	0.68 \pm 0.02	30.0 \pm 2.0	5.7	0.98 \pm 0.02	17 \pm 31	1.2/239
11 ⁴	56591.375	56591.382	0.61	0.23 \pm 0.02	2.2	0.71 \pm 0.03	28.1 \pm 2.1	6.1	0.98 \pm 0.02	27 $^{+25}_{-46}$	1.1/253
12	56592.450	56592.462	1.05	0.21 \pm 0.01	2.2	0.71 \pm 0.02	26.2 $^{+1.8}_{-1.0}$	6.0	0.95 \pm 0.01	39 \pm 20	1.2/324
13	56592.058	56592.062	0.34	0.22 \pm 0.05	2.2	0.70 \pm 0.04	31.7 $^{+6.3}_{-7.2}$	3.6	1.02 \pm 0.02	0 $^{+56}_{-72}$	0.98/164
14	56594.177	56594.994	1.48	0.22 \pm 0.02	2.2	0.72 \pm 0.02	28.9 \pm 1.5	5.9	0.95 \pm 0.01	33 $^{+22}_{-27}$	1.2/330
15	56595.912	56595.921	0.74	0.24 \pm 0.03	2.2	0.70 \pm 0.02	30.2 $^{+3.1}_{-2.2}$	6.2	0.98 \pm 0.01	0 $^{+30}_{-40}$	1.1/302
16	56596.120	56596.132	0.98	0.20 \pm 0.01	2.2	0.71 \pm 0.01	27.3 \pm 1.3	5.6	0.95 \pm 0.01	40 $^{+16}_{-23}$	1.1/308
17	56597.180	56597.189	0.76	0.22 \pm 0.01	2.2	0.67 \pm 0.02	28.6 $^{+2.8}_{-1.7}$	5.8	0.95 \pm 0.01	40 $^{+18}_{-23}$	1.1/258
18	56598.915	56598.927	1.05	0.24 \pm 0.04	2.2	0.68 \pm 0.03	28.4 $^{+4.0}_{-2.0}$	4.6	1.01 \pm 0.01	0 $^{+32}_{-50}$	0.98/294
19	56599.453	56599.467	1.16	0.20 \pm 0.01	2.2	0.71 \pm 0.02	25.5 $^{+3.7}_{-1.4}$	4.9	0.95 \pm 0.02	51 $^{+26}_{-21}$	1.2/299
20	56600.987	56601.000	1.15	0.21 \pm 0.01	2.2	0.68 \pm 0.02	24.9 \pm 1.7	4.9	0.95 \pm 0.01	38 $^{+18}_{-22}$	1.1/273
21	56601.384	56601.393	0.80	0.21 \pm 0.02	2.2	0.67 \pm 0.02	25.3 \pm 1.6	4.7	0.96 \pm 0.01	33 $^{+18}_{-22}$	1.1/264
22	56604.994	56604.997	0.19	0.3 \pm 0.1	2.2	0.6 \pm 0.1	38.5 $^{+12.3}_{-15.3}$	4.8	1.0 \pm 0.1	0 $^{+194}_{-93}$	1.1/108
23	56605.928	56605.940	1.05	0.21 \pm 0.01	2.2	0.62 \pm 0.02	26.9 \pm 2.2	4.6	0.95 \pm 0.01	41 $^{+22}_{-18}$	1.3/258
24	56606.121	56606.130	0.75	0.25 \pm 0.03	2.2	0.61 \pm 0.02	29.4 \pm 2.8	4.6	0.96 \pm 0.02	12 $^{+33}_{-48}$	1.0/251
25	56607.670	56607.681	0.95	0.22 \pm 0.02	2.2	0.64 \pm 0.02	31.1 $^{+3.2}_{-1.6}$	5.7	0.96 \pm 0.01	36 $^{+18}_{-29}$	1.2/253

Notes. ¹XRT observation ID as in Table 1. ²The radius of the disk is estimated as described in the notes of Table 1. ³The value of Γ was fixed to that of the averaged broad-band spectrum (Fig. 3). ⁴The observation was divided into two snapshots that we analyzed separately. ⁵As in Table 1.




Fig. 4. Averaged PSD of MAXI J1828-249 obtained from all available XRT data. The PSD here is rebinned with a logarithmic step of 1.25. The white noise contribution has been subtracted. The red solid line represents the best fit obtained with a zero-centered Lorentzian (see text for details).




Fig. 5. Fractional rms of MAXI J1828-249 (4-50 mHz frequency range) measured from the XRT data. The left panel shows the rms as function of time; in the right panel the average XRT count-rate of each pointing is plotted as a function of the fractional rms.

spectral transition from a hard to a softer state about one day later (Negoro et al. 2013). Such transitions are typically observed in the so-called black hole candidates (BHC), which suggests that MAXI J1828-249 is associated with this class of objects. According to the standard scenario (Fender et al. 2004; Homan & Belloni 2005; Belloni 2010), BHC sources are known to evolve during their outburst along a q-shaped track in the hardness-intensity diagram. The outburst starts in the low-hard spectral state (LHS), which is characterized by a power-law shaped X-ray spectrum with $\Gamma \sim 1.6 - 1.7$ and a cut-off at the higher energies $E_{cut} \sim 100$ keV. Radio emission arises in this state due to the synchrotron radiation of a steady jet. In the following phases of the outburst, the X-ray and radio luminosities both increase until the source reaches the high-soft state (HSS), characterized by a prominent thermal emission from the accretion disk and a marginal power-law tail. Radio emission at this state is no longer observed, most likely because of the suppression of the jet. The time variability of the source is also significantly different in the two states, being generally more pronounced in the LHS with rms values of up to $\sim 30\%$ and characterized by the quasi-periodical oscillations (QPOs) in the HSS in combination with a strongly suppressed rms (see, e.g., Belloni 2010, for a recent review). In addition to these two main states, at least two intermediate hard and soft states were identified. In particular, BHCs in the so-called intermediate soft state still show a significant hard component (extending to hundreds of keV) though they are no longer significantly detected in the radio domain and present both a prominent soft spectral component and a limited rms. Not all transient BHCs in outburst go through a complete q-track. So far, a limited number of objects were observed that reach the hard intermediate state during outburst, but then return to the LHS instead of moving to HSS (see, e.g., Capitanio et al. 2009; Ferrigno et al. 2012).

The observational campaign we presented in this paper started about 1.8 days after the discovery, that is, immediately after the possible spectral transition reported by Negoro et al. (2013). The broad-band energy spectrum of the source as mea-

sured by *Swift* and *INTEGRAL* comprised a thermal component that we associated with the emission from a multi-temperature black-body disk, and a hard power-law extending with no measurable break up to 200 keV. Assuming a distance to the source of 8 kpc, its peak luminosity is $\sim 10^{38}$ erg/s. During the entire observational campaign MAXIJ1828-249 did not show significant spectral evolution. Our observational results suggest that MAXIJ1828-249 is a transient BHC that evolved rapidly in the first two days of the outburst from the LHS to the soft intermediate state and remained in this state during the entire observational campaign presented in this work. This conclusion is supported by both the spectral and timing analysis⁴ reported in Sect. 4 and the lack of significant radio emission already two days after the onset of the outburst, as mentioned in Sect. 1. The light curve of the source based on MAXI publicly available data also supports this scenario and suggests that the source is now fading into quiescence. This means that MAXIJ1828-249 might represent another example of transient BHC "failed" outburst, as mentioned above. Future observations in mid-2014 (e.g. with *Swift*/XRT) will be able to confirm/reject this conclusion.

Note added to proofs. After this paper was accepted for publication, new *Swift* data collected on 2014 February 14 found that the source is back in a faint hard state (power-law photon index ~ 1.7 and 0.5-10 keV flux of 4×10^{-11} erg/cm²/s; Tomsick & Corbel (2014)). Radio observations performed on 2014 February 16 also detected significant radio emission from the source (preliminary flux densities 1.38 ± 0.05 mJy at 5.5 GHz and 1.28 ± 0.06 mJy at 9 GHz; Corbel et al. (2014)), which supports our conclusions in Sect. 5.

Acknowledgments

EF acknowledges support by grant NSH-6137.2014.2, RFBR 13-02-00741 and grant 20FI20_135263 of the National Swiss Fond.

References

- Belloni, T. M. 2010, in *Lecture Notes in Physics*, Berlin Springer Verlag, Vol. 794, *Lecture Notes in Physics*, Berlin Springer Verlag, ed. T. Belloni, 53–+
 Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
 Capitanio, F., Belloni, T., Del Santo, M., & Ubertini, P. 2009, *MNRAS*, 398, 1194
 Corbel, S., Tomsick, J. A., & Tzioumis, T. 2014, *The Astronomer's Telegram*, 5911, 1
 Courvoisier, T., Walter, R., Beckmann, V., et al. 2003, *A&A*, 411, L53
 Dickey, J. M. & Lockman, F. J. 1990, *ARA&A*, 28, 215
 Fender, R. P., Belloni, T. M., & Gallo, E. 2004, *MNRAS*, 355, 1105
 Ferrigno, C., Bozzo, E., Del Santo, M., & Capitanio, F. 2012, *A&A*, 537, L7
 Filippova, E., Kuulkers, E., Sanchez-Fernandez, C., et al. 2013, *The Astronomer's Telegram*, 5476, 1
 Homan, J. & Belloni, T. 2005, *Ap&SS*, 300, 107
 Kennea, J. A., Linares, M., Krimm, H. A., et al. 2013a, *The Astronomer's Telegram*, 5478, 1
 Kennea, J. A., Linares, M., Krimm, H. A., et al. 2013b, *The Astronomer's Telegram*, 5479, 1
 Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141
 Lund, N., Budtz-Jørgensen, C., Westergaard, N. J., et al. 2003, *A&A*, 411, L231
 Miller-Jones, J. C. A., Russell, T. D., Sivakoff, G. R., & Curran, P. A. 2013, *The Astronomer's Telegram*, 5484, 1
 Nakahira, S., Tomida, H., Negoro, H., et al. 2013, *The Astronomer's Telegram*, 5474, 1

- Negoro, H., Sugizaki, M., Mihara, T., et al. 2013, *The Astronomer's Telegram*, 5483, 1
 Rau, A., Tanga, M., & Greiner, J. 2013, *The Astronomer's Telegram*, 5482, 1
 Romano, P., Campana, S., Chincarini, G., et al. 2006, *A&A*, 456, 917
 Tomsick, J. A. & Corbel, S. 2014, *The Astronomer's Telegram*, 5886, 1
 Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131

⁴ Note that the range of frequencies we could investigate in the case of MAXIJ1828-249 based on the available X-ray data is slightly lower (0.001-0.5 Hz) compared to that in which typical QPOs for this state are observed (0.01 - 50 Hz). However, the measured rms level in the XRT data is fully compatible with that commonly found in the soft intermediate state.